

C11

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/415, 31/155		A1	(11) International Publication Number: WO 95/19772
			(43) International Publication Date: 27 July 1995 (27.07.95)
(21) International Application Number: PCT/US95/00504 (22) International Filing Date: 18 January 1995 (18.01.95) (30) Priority Data: 185,079 20 January 1994 (20.01.94) US (60) Parent Application or Grant (63) Related by Continuation US 185,079 (CON) Filed on 20 January 1994 (20.01.94)		(74) Agent: SIBLEY, Kenneth, D.; Bell, Seltzer, Park & Gibson, P.O. Drawer 34009, Charlotte, NC 28234 (US). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).	
(71) Applicant (for all designated States except US): THE UNI- VERSITY OF NORTH CAROLINA AT CHAPEL HILL [US/US]; 300 Bynum Hall, Campus Box 4100, Chapel Hill, NC 27599-4100 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DYKSTRA, Christine, C. [US/US]; 203 Bennington Drive, Chapel Hill, NC 27516 (US). SWANSTROM, Ronald, I. [US/US]; 7021 Knotty Pine Drive, Chapel Hill, NC 27514 (US). TIDWELL, Richard, R. [US/US]; 1806 The Oaks, Chapel Hill, NC 27514 (US).		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: USE OF BIS(AMIDINOBENZIMIDAZOLES) IN THE MANUFACTURE OF MEDICAMENT FOR INHIBITING RETRO- VIRAL INTEGRASE			
(57) Abstract Disclosed is a method of inhibiting retroviral integrase in a subject in need of such treatment. The method comprises administering to the subject an effective retroviral integrase inhibiting amount of a bis-benzimidazole compound such as bis[5-amidino-2-benzimidazolyl]methane, 1,2-bis[5-amidino-2-benzimidazolyl]ethane, 1,2-bis[5-amidino-2-benzimidazolyl]ethene or a pharmaceutically acceptable salt thereof. A method for combatting retroviral infections is also disclosed.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

USE OF BIS(AMIDINO BENZIMIDAZOLES) IN THE MANUFACTURE OF MEDICAMENT FOR
INHIBITING RETROVIRAL INTEGRASE

Field of the Invention

The present invention relates to methods of inhibiting retroviral integrases with dicationic bis-benzimidazoles.

5 Background of the Invention

The genome of the typical retrovirus encodes three primary enzymes that mediate the virus replication cycle. Reverse transcriptase converts the viral RNA genome into a double stranded DNA. Integrase
10 nonspecifically inserts this DNA copy into the host cell genome, and protease cleaves viral structural and nonstructural proteins into their mature forms.

An essential step of the retroviral life cycle is the integration of its double-stranded DNA
15 copy into the host genome. H. Sakai et al, *J. Virol.* 67, 1169-74 (1993). This process requires highly conserved sequence recognition and cleaving steps. For this reason, a therapeutic agent that can interrupt this process should be an effective and specific
20 antiviral agent. A protein at the C-terminus of the viral polymerase, integrase (IN), is the only viral protein required for this process. R. LaFemina et al., *J. Virol.* 66, 7414-7419 (1992).

A. Fesen et al., *Proc. Natl. Acad. Sci. USA*
25 90, 2399-2403 (1993) discuss investigations using an *in vitro* integrase assay of a variety of chemicals as potential human immunodeficiency virus type I (HIV-1) integrase inhibitors. The article reports several


-2-

topoisomerase inhibitors, such as doxorubicin, mitoxantrose, ellipticines and quercetin as potent inhibitors. While some topoisomerase inhibitors were excellent anti-integrase agents, no correlation was
 5 observed with antiviral effects. This is believed to be at least partially due to the fact that a number of topoisomerase inhibitors have severe cytotoxic effects, depending upon their mechanism of inhibition.

R. LaFemina et al., *J. Virology* 56, 7414-7419
 10 (1992) reports studies assessing the usefulness of the integrase enzyme as a target for specific HIV-1 anti-viral therapeutic agents by determining its absolute requirement for productive HIV-1 infection. The article reports the results of the introduction of
 15 specific amino acid substitution into recombinant integrase and assesses the ability of the mutant proteins to properly mediate specific and non-specific cleavage as well as integration.

Summary of the Invention

20 A first aspect of the present invention is a method of inhibiting retroviral integrase (e.g. *in vitro*, or in a subject in need of such treatment). The method comprises administering to the subject or contacting to the retroviral integrase an effective
 25 retroviral integrase-inhibiting amount of a compound according to Formula (I):



wherein:

R_1 and R_2 are each independently selected from the group consisting of H or lower alkyl, or R_1 and R_2 together represent $-(CH_2)_m-$
 30 wherein m is from two to four;
 R_3 is H or lower alkyl; and

-3-

X is C1 to C2 saturated or unsaturated alkyl containing up to one double bond (e.g., $-(CH_2)_n$ wherein n is from 1-2, which is unsubstituted or substituted from 1 to 2
5 times with lower alkyl, and which is saturated or unsaturated and contains up to one double bond);

or a pharmaceutically acceptable salt thereof.

In a preferred embodiment of the invention, R_1
10 and R_2 are each H; R_2 is H or lower alkyl; and x is selected from the group consisting of $-CH_2-CH_2-$ and $-CH=CH-$ and any of the foregoing substituted from 1 to 2 times with lower alkyl; and the pharmaceutically acceptable salts thereof. Currently preferred
15 compounds are bis[5-amidino-2-benzimidazolyl]methane; 1,2-bis[5-amidino-2-benzimidazolyl]ethane; 1,2-bis[5-amidino-2-benzimidazolyl]ethene; 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethene, or pharmaceutically acceptable salts thereof.

20 A second aspect of the present invention is a method of combatting a retroviral infection (e.g. in vitro, or in a subject in need of such treatment). The method comprises administering to the subject or contacting to the retrovirus an effective retroviral
25 infection combatting amount of a compound according to Formula (I) as given above or a pharmaceutically acceptable salt thereof. Currently preferred compounds are bis[5-amidino-2-benzimidazolyl]methane; 1,2-bis[5-amidino-2-benzimidazolyl]ethane; 1,2-bis[5-amidino-2-benzimidazolyl]ethene; 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethene, or pharmaceutically acceptable
30 salts thereof.

Compounds of formula I, or there
pharmaceutically acceptable salts, may be included in a
35 therapeutically effective amount in a pharmaceutically acceptable carrier to provide pharmaceutical formulations, with the therapeutically effective amount

-4-

being effective to carry out the methods set forth above.

Further aspects of the present invention include the use of compounds of Formula (I) above and
5 their pharmaceutically acceptable salts for the preparation of a medicament for inhibiting retroviral integrase, or for combatting a retroviral infection.

The foregoing and other objects and aspects of the present invention are explained in detail in the
10 specification set forth hereinbelow.

Brief Description of the Drawings

Figure 1 shows the plating efficiency of CEM cells, a human T-lymphoma cell line (A.H. Kaplan et al, *J. Virol.* 67, 4050-5 (1993)), treated with various
15 concentrations of 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethene (Compound B) ($30\mu\text{M}$, $100\mu\text{M}$, and $300\mu\text{M}$), the x-axis representing exposure time in days and the y-axis representing viable cells per milliliter (ml); and

20 Figure 2 shows the growth rate of Magic cells (J. Kimpton et al, *J. Virol.* 66, 2232-2239 (1992)) in the presence of various concentrations of 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethene (Compound B) ($10\mu\text{M}$, $30\mu\text{M}$, and $100\mu\text{M}$), the x-axis representing
25 exposure time in days and the y-axis representing the number of cells.

Detailed Description of the Invention

The term "lower alkyl," as used herein, refers to C1 to C4 linear or branched alkyl, such as
30 methyl, ethyl, propyl, butyl, isopropyl, sec-butyl, and tert-butyl. Methyl and ethyl are preferred.

Subjects to be treated by the methods of the present invention are animal subjects, typically vertebrates, including both mammalian (e.g., human,
35 cat, dog, cow, horse, sheep, pig, monkey, ape, etc.) and avian subjects (e.g., chicken, turkey, duck, goose, quail, pheasant, etc.).

-5-

The invention applies generally to retroviruses, i.e., the entire Retroviridae virus family. The family encompasses all viruses containing an RNA genome and an RNA-dependent DNA polymerase (reverse transcriptase) enzymatic activity. The family is divided into three subfamilies: (1) Oncovirinae, including all the oncogenic members and many closely related nononcogenic viruses; (2) Lentivirinae, the "slow" viruses, such as visna virus; and (3) Spumavirinae, the "foamy" viruses that induce persistent infections without any clinical disease. Retroviruses of interest include human retroviruses, such as immunodeficiency virus type 1 (HIV-1), avian retroviruses, such as avian sarcoma and leukosis viruses of chickens (ASLVs), endogenous viruses of certain pheasant and quail species, reticuloendotheliosis virus of turkeys and related viruses of ducks and chickens, and lymphoproliferate disease virus of turkeys; feline C-type retroviruses, including feline leukemia virus (FeLV) and feline sarcoma virus (FeSV) and endogenous retroviruses (RD114 and CCC isolates); mink C-type retroviruses, including mink leukemia virus (MiLV); porcine C-type retroviruses; horse C-type retroviruses, including the equine infectious anemia virus (EIAV); bovine C-type retrovirus, including enzootic bovine leukosis or lymphosarcoma; sheep C-type retroviruses; and primate retroviruses, including prosimian C-type retroviruses, Simian sarcoma and gibbon ape leukemia C-type retroviruses, baboon C-type retroviruses, macaque C-type retroviruses, owl monkey C-type retroviruses, Colobus monkey C-type retroviruses, Mason-Pfizer monkey D-type retrovirus, Langur D-type retrovirus and squirrel monkey D-type retrovirus. See N. Teich, *Taxonomy of Retroviruses in Molecular Biology of Tumor Viruses*, R. Weiss, N. Teich, H. Varmus, and J. Coffin,

-6-

Eds., Cold Spring Harbor Laboratory, New York (2d ed. 1984), pps. 26-207.

As noted above, the present invention provides pharmaceutical formulations comprising the
5 aforementioned compounds of Formula I, or
pharmaceutically acceptable salts thereof, in
pharmaceutically acceptable carriers, preferably for
aerosol, oral, and parenteral administration as
discussed in greater detail below. Also, the present
10 invention provides such compounds or salts thereof
which have been lyophilized and which may be
reconstituted to form pharmaceutically acceptable
formulations for administration, as by intravenous or
intramuscular injection.

15 The therapeutically effective dosage of any
specific compound, the use of which is in the scope of
the present invention, will vary somewhat from compound
to compound, patient to patient, and will depend upon
the condition of the patient and the route of delivery.
20 As a general proposition, a dosage from about 0.1 to
about 20 mg/kg will have therapeutic efficacy, with
still higher dosages potentially being employed for
oral and/or aerosol administration. Toxicity concerns
at the higher level may restrict intravenous dosages to
25 a lower level such as up to about 10 mg/kg, all weights
being calculated based upon the weight of the active
base, including the cases where salt is employed.
Typically a dosage from about 0.56 mg/kg to about 5
mg/kg will be employed. In certain circumstances,
30 higher or lower doses may be also appropriate. The
duration of the treatment may be once per day for a
period of from two to three weeks and may continue for
period of months or even years, for example, in
treating chronic conditions. Lower doses given less
35 frequently can be used to prevent or reduce the
incidence of recurrence of the infection. The daily
dose can be administered either by a single dose in the

-7-

form of an individual dosage unit or several smaller dosage units or by multiple administration of subdivided dosages at certain intervals.

Compounds of Formula (I) may be administered
5 per se or in the form of a pharmaceutically acceptable salt. When used in medicine, the salts of the compounds of Formula (I) should be both pharmacologically and pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently
10 be used to prepare the free active compound or pharmaceutically acceptable salts thereof and are not excluded from the scope of this invention. Such pharmacologically and pharmaceutically acceptable salts include, but are not limited to, those prepared from
15 the following acids: hydrochloric, lactate, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluenesulfonic, tartaric, citric, methanesulfonic, formic, malonic, succinic, naphthalene-2-sulfonic and benzenesulfonic. Also,
20 pharmaceutically acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts of the carboxylic acid group. Thus, the present invention also provides pharmaceutical formulations, both for veterinary and
25 for human medical use, which comprise the retroviral integrase inhibiting agent together with one or more pharmaceutically acceptable carriers thereof and optionally any other therapeutic ingredients. The carrier(s) must be pharmaceutically acceptable in the
30 sense of being compatible with the other ingredients of the formulation and not unduly deleterious to the recipient thereof.

The formulations include those suitable for oral, rectal, topical, nasal, ophthalmic or parenteral
35 (including subcutaneous, intramuscular and intravenous) administration. Formulations suitable for aerosol, oral and parenteral administration are preferred.

-8-

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active
5 compound into association with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided
10 solid carrier, or both, and then, if necessary, shaping the product into desired formulations.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, tablets or
15 lozenges, each containing a predetermined amount of the integrase inhibiting agent as a powder or granules; or a suspension in an aqueous liquor or non-aqueous liquid such as a syrup, an elixir, an emulsion or a draught.

A tablet may be made by compression or
20 molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine, with the active compound being in a free-flowing form such as a powder or granules which is optionally mixed with a binder,
25 disintegrant, lubricant, inert diluent, surface active agent or dispersing agent. Molded tables comprised of a mixture of the powdered active compound with a suitable carrier may be made by molding in a suitable machine.

30 A syrup may be made by adding the active compound to a concentrated aqueous solution of a sugar, for example sucrose, to which may also be added any accessory ingredient(s). Such accessory ingredient(s) may include flavorings, suitable preservatives, an
35 agent to retard crystallization of the sugar, and an agent to increase the solubility of any other

- 9 -

ingredient, such as a polyhydric alcohol, for example glycerol or sorbitol.

Formulations suitable for parenteral administration conveniently comprise a sterile aqueous
5 preparation of the active compound, which is preferably isotonic with the blood of the recipient and pyrogen-free.

Nasal spray formulations comprise purified aqueous solutions of the active compound with
10 preservative agents and isotonic agents. Such formulations are preferably adjusted to a pH and isotonic state compatible with the nasal mucous membranes.

Formulations for rectal administration may be
15 presented as a suppository with a suitable carrier such as cocoa butter, or hydrogenated fats or hydrogenated fatty carboxylic acids.

Ophthalmic formulations are prepared by a similar method to the nasal spray, except that the pH
20 and isotonic factors are preferably adjusted to match that of the eye.

Topical formulations comprise the active compound dissolved or suspended in one or more media such as mineral oil, petroleum, polyhydroxy alcohols or
25 other bases used for topical pharmaceutical formulations. The addition of other accessory ingredients, vide infra, may be desirable.

In addition to the aforementioned ingredients, the formulations of this invention may
30 further include one or more accessory ingredient(s) selected from diluents, buffers, flavoring agents, binders, disintegrants, surface active agents, thickeners, lubricants, preservatives (including antioxidants) and the like.

35 In accordance with the preferred embodiments of the present method, a compound of Formula I, or a pharmaceutically acceptable salt thereof, may be

-10-

administered orally or through inhalation as a solid, or may be administered intramuscularly or intravenously as a solution, suspension, or emulsion. Alternatively, the compound or salt may also be administered by

5 inhalation, intravenously or intramuscularly as a liposomal suspension. When administered through inhalation the compound or salt should be in the form of a plurality of solid particles or droplets having a particle size from about 0.5 to about 5 microns,

10 preferably from about 1 to about 2 microns.

The present invention also provides pharmaceutical compositions suitable for intravenous or intramuscular injection. The pharmaceutical compositions comprise a compound of Formula I, or a

15 pharmaceutically acceptable salt thereof, in any pharmaceutically acceptable carrier. If a solution is desired, water is the carrier of choice with respect to water-soluble compounds or salts, an organic vehicle, such as glycerol, propylene, glycol, polyethylene

20 glycol, or mixtures thereof, may be suitable. In the latter instance, the organic vehicle may contain a substantial amount of water. The solution in either instance may then be sterilized in any suitable manner, preferably by filtration through a 0.22 micron filter.

25 Subsequent to sterilization, the solution may be filled into appropriate receptacles, such as depyrogenated glass vials. Of course, the filling should be done by an aseptic method. Sterilized closures may then be placed on the vials and, if desired, the vial contents

30 may be lyophilized.

In addition to compounds of Formula I or their salts, the pharmaceutical compositions may contain other additives, such pH adjusting additives. In particular, useful pH adjusting agents include acids

35 or bases or buffers, such a sodium lactate, sodium acetate, or sodium gluconate. Further, the compositions may contain microbial preservatives.

-11-

Useful microbial preservatives include methylparaben, propylparaben, and benzyl alcohol. The microbial preservative is typically employed when the formulation is placed in a vial designed for multidose use. Of course, as indicated, the pharmaceutical compositions of the present invention may be lyophilized using techniques well known in the art.

In yet another aspect of the present invention, there is provide an injectable, stable, sterile composition comprising a compound of Formula I, or a salt thereof, in a unit dosage form in a sealed container. The compound or salt is provided in the form of a lyophilizate which is capable of being reconstituted with a suitable pharmaceutically acceptable carrier to form a liquid composition suitable for injection thereof into the subject. The unit dosage form typically comprises from about 10 mg to about 10 grams of the compound or salt. When the compound or salt is substantially water-insoluble, a sufficient amount of emulsifying agent which is physiologically acceptable may be employed in sufficient quantity to emulsify the compound or salt in an aqueous carrier. One such useful emulsifying agent is phosphatidyl choline.

Other pharmaceutical compositions may be prepared from the water-insoluble compounds of Formula I, or salts thereof, such as aqueous base emulsions. In such an instance, the composition will contain a sufficient amount of pharmaceutically acceptable emulsifying agent to emulsify the desired amount of the compound of Formula I or salt thereof. Particularly useful emulsifying agents include phosphatidyl cholines, and lecithin.

Further, the present invention provides liposomal formulations of the compounds of Formula I and salts thereof. The technology for forming liposomal suspensions is well known in the art. When

-12-

the compound of Formula I or salt thereof is an aqueous-soluble salt, using conventional liposome technology, the same may be incorporated into lipid vesicles. In such an instance, due to the water solubility of the compound or salt, the compound or salt will be substantially entrained within the hydrophilic center or core of the liposomes. The lipid layer employed may be of any conventional composition and may either contain cholesterol or may be cholesterol-free. When the compound or salt of interest is water-insoluble, again employing conventional liposome formation technology, the salt may be substantially entrained within the hydrophobic lipid bilayer which forms the structure of the liposome. In either instance, the liposomes which are produced may be reduced in size, as through the use of standard sonication and homogenization techniques.

Of course, the liposomal formulations containing the compounds of Formula I or salts thereof, may be lyophilized to produce a lyophilizate which may be reconstituted with a pharmaceutically acceptable carrier, such as water, to regenerate a liposomal suspension.

Pharmaceutical formulations are also provided which are suitable for administration as an aerosol, by inhalation. These formulations comprise a solution or suspension of the desired compound of Formula I or a salt thereof or a plurality of solid particles of the compound or salt. The desired formulation may be placed in a small chamber and nebulized. Nebulization may be accomplished by compressed air or by ultrasonic energy to form a plurality of liquid droplets or solid particles comprising the compounds of salts. The liquid droplets or solid particles should have a particle size in the range of about 0.5 to about 5 microns. The solid particles can be obtained by processing the solid compound of Formula I, or a salt

-13-

thereof, in any appropriate manner known in the art, such as by micronization. Most preferably, the size of the solid particles or droplets will be from about 1 to about 2 microns. In this respect, commercial

5 nebulizers are available to achieve this purpose. Preferably, when the pharmaceutical formulation suitable for administration as an aerosol is in the form of a liquid, the formulation will comprise a water-soluble compound of Formula I or a salt thereof,
10 in a carrier which comprises water. A surfactant may be present which lowers the surface tension of the formulation sufficiently to result in the formation of droplets within the desired size range when subjected to nebulization.

15 As indicated, the present invention provides both water-soluble and water-insoluble compounds and salts. As used in the present specification, the term "water-soluble" is meant to define any composition which is soluble in water in an amount of about 50
20 mg/mL, or greater. Also, as used in the present specification, the term "water-insoluble" is meant to define any composition which has solubility in water of less than about 20 mg/mL. For certain applications, water soluble compounds or salts may be desirable
25 whereas for other applications water-insoluble compounds or salts likewise may be desirable.

Examples of compounds exemplary of Formula (I) above include, but are not limited to:

- (A) 1,2-bis[5-amidino-2-
30 benzimidazolyl]ethene;
- (B) 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethene;
- (C) bis[5-(2-imidazolyl)-2-benzimidazolyl]
methane;
- 35 (D) bis[5-amidino-2-benzimidazolyl] methane;
- (E) (5-Amidino-2-benzimidazolyl)-(5'-(2-imidazolyl)-2'-benzimidazolyl)methane;

-14-

(AA) 1,2-bis[5-amidino-2-benzimidazolyl]ethane;

(BB) 1,2-bis[5-(2-imidazolyl)-2-benzimidazolyl]ethane;

5 (CC) 1,2-bis[5-(2-imidazolyl)-2-benzimidazolyl]ethene;

(DD) 1-(5-Amidino-2-benzimidazolyl)-(5'-(2-imidazolyl)-2'-benzimidazolyl)ethane;

(EE) 1-(5-Amidino-2-benzimidazolyl)-(5'-(2-imidazolyl)-2'-benzimidazolyl)ethene;

10 (FF) bis[5-isopropylamidino-2-benzimidazolyl]methane; and

(GG) 1,2-bis[5-isopropylamidino-2-benzimidazolyl]ethane.

15 Additional compounds illustrative of Formula (I) analogous to the foregoing can be produced by substituting a 2-pyrimidyl group for the 2-imidazolyl group.

Compounds employed in carrying out the present invention are either known or may be prepared in accordance with techniques known to those skilled in the art (see, e.g., U.S. Patent No. 4,933,347, the entire disclosure of which is hereby incorporated by reference), particularly in light of the disclosure set forth below.

As indicated, the compounds used in the present invention may be present as pharmaceutically acceptable salts. Such salts include the gluconate, lactate, acetate, tartrate, citrate, phosphate, borate, nitrate, sulfate, and hydrochloride salts.

The salts of the present invention may be prepared, in general, by reacting two equivalents of the amidine base compound with the desired acid, in solution. After the reaction is complete, the salts are crystallized from solution by the addition of an appropriate amount of solvent in which the salt is insoluble.

-15-

The present invention will be further illustrated by the following non-limiting examples, in which "g" means grams, "mg" means milligrams, " μ g" means micrograms, "mmol" means millimoles, "h" means hours, "ml" means milliliter, "M" means molar, "mM" means millimolar, "UV" means ultraviolet, "HCl" means hydrogen chloride, "NaCl" means sodium chloride, "EDTA" means ethylenediaminetetraacetic acid, "MP" means melting point, and " $^{\circ}$ C" means degrees Celcius.


EXAMPLE 1

Preparation of Compounds


The structures of the compounds used in the inhibition studies described below are given in Table 1 below. All were synthesized by previously published methods. See R. Tidwell et al, *Antimicrob. Agents and Chemother.* 38, 000-000 (1993); T. Fairley, *J. Med. Chem.* 36, 1746-1753 (1993).

TABLE 1

Structures of Integrase Inhibiting Compounds



Compound	R ₁	R ₂	X
A	Am	Am	-C=C-
B	IsopropAm	IsopropAm	-C=C-
C	Im	Im	CH ₂
D	Am	Am	CH ₂
E	Am	Im	CH ₂



-16-

EXAMPLE 2

Preparation and Assay of Integrase Protein

Purification of Integrase Protein. Integrase was overproduced in *E. coli* from an inducible plasmid containing the integrase sequence under control of a *lacI*-regulated T7 polymerase using the integrase coding domain from the HXB2 clone of HIV-1 DNA (L. Ratner et al, *AIDS Res. Hum. Retroviruses* 3, 57-69 (1987)), pT7f11-IN. A one liter culture at an O.D. of 0.6 was induced by the addition of IPTG to 0.5 mM. After three hours, the cells were harvested by centrifugation and the pellets were stored at -70°C.

The integrase protein was purified by a modification of the method of Sherman and Fyfe (P. Sherman et al, *Proc. Natl. Acad. Sci. USA* 87, 5119-23 (1990)). The cells were lysed by thawing in a buffer (50 mM Tris-HCl pH 7.5, 5 mM dithiothreitol, 1 mM EDTA, 1 mg/ml lysozyme) on ice at 6 ml/g bacterial pellet for 30 minutes, and then incubating at 37°C for 5 minutes. The lysate was centrifuged at 12,000 X g for one hour. The pellets were resuspended in 50 mM Tris-HCl, pH 7.5, 1 mM dithiothreitol, 1 mM EDTA, 1 M NaCl (4 ml/g original bacteria). The homogenate was stirred for 30 minutes at 4°C and recentrifuged for 30 minutes at 12,000 X g.

The supernatant was made 0.8 M in ammonium sulfate by the slow addition of powder with stirring over a 30 minute period. The extract was then centrifuged to remove any precipitate and applied to a phenyl sepharose column. After a 50 ml wash with high salt buffer (50 mM Tris-HCl, 1 mM EDTA, 5 mM DTT, 2 M NaCl), the protein was eluted with a gradient from high salt to 0 salt buffer containing 10% (weight/vol) glycerol and further purified through a G75 Sephadex column to remove background nuclease. A one liter culture generates enough integrase activity to perform over a thousand drug inhibition assays.

-17-

Assays of Integrase Activity. The enzyme purification steps were monitored by an endonuclease/nicking assay and western blots utilizing a monoclonal antibody to integrase generated by W. Osheroff and R. Swanstrom at UNC - Chapel Hill. Supercoiled pBluescriptKS+II (0.3 μ g) was incubated with column fractions in a buffer containing 20 mM Tris-HCl, pH 8.0, 5 mM 2-mercaptoethanol, and 2 mM $MnCl_2$ for 30 minutes at 37°C. The reactions were stopped by the addition of SDS to 1% and cleavage of the DNA substrate is assessed by electrophoresis through a 0.8% agarose gel stained with ethidium bromide and photographed under UV illumination.

Cleavage of specific sites is assessed as described in Sherman et al., *supra*, except that the assay buffer was the same as that reported by Chow et al, *Science* 255, 723-6 (1992). Single 20 mer oligonucleotides corresponding to either the U5 or U3 ends of HIV-1 are end-labeled with ^{32}P , annealed with its complement, purified, and used under the same conditions as described above for the endonuclease assay. Reaction products are denatured with formamide, electrophoresed through 20% denaturing polyacrylamide gels and visualized by autoradiography. Both cleavage and ligation activity can be assessed from one gel. To assess the ligation activity alone, "precleaved" substrate that had the -2 cleavage artificially produced by using a radiolabelled 18 mer oligonucleotide was also used for some experiments.

30

EXAMPLE 3

Virus and Cell Cultures

Cell lines used were CEM cells, a human T-cell lymphoma cell line, A. Kaplan et al, *J. Virol.* 67, 4050-5 (1993), and Magic cells, J. Kimpton et al, *J. Virol.* 66, 2232-2239 (1992). The CEM cells were grown in RPMI-1640 medium supplemented with 5% FCS. The Magic Cells, a HELA derivative, were grown in DMEM/H

-18-

supplemented with 5% FCS, G418 (20 mg/ml) and Hygromycin (10 mg/ml). The HIV isolate was strain HXB2, originally from Lee Ratner in the laboratory of Robert Gallo at the National Institutes of Health.

5

EXAMPLE 4**Toxicity Assays**

Three different toxicity tests were performed on the cell lines used to measure virus infectivity.

The initial test for toxicity utilized the

10 XTT Assay as described by Weislow et al, *J. Natl. Cancer Inst.* **81**, 577-586 (1989). This is the standard assay used originally to measure cell toxicity by potential reverse transcriptase inhibitors. Briefly, cells were grown to cells/ml and drug dilutions were

15 added to the medium. After two days incubation, XTT reagent is added and incubation is continued for 4h at 37°C. Following incubation, the plates are read at 450 minus 650 nm (the 650 value is the background value which is automatically subtracted) with controls of

20 media + XTT reagent without cells, and cells + media without reagent. Also a control of cells + media + XTT reagent was run for each plate. Medium without phenol red was employed to minimize background color as the XTT reagent goes from colorless (unreduced) to orange

25 (reduced). The XTT reagent was freshly prepared on the day of assay as follows: 1 mg/ml of XTT in 0.01 M phenazine methosulfate. The phenazine methosulfate solution was prepared ahead of time and stored at 4°C in a dark bottle. The XTT reagent is added to the

30 microtiter wells at 24 ul per 100 ul of medium. The O.D. was measured on a Vmax plate reader from Molecular Devices Co. with data reduction. Results are expressed as percent of untreated controls. The least toxic compounds were Compounds A, B, and F with toxicities

35 values at 500 μ M or greater.

Next, a plating efficiency test measured the ability of the cell to grow after incubation with drug

-19-

for a number of days. Magic cells were grown from an initial cell concentration of 0.8×10^4 with or without various concentrations of test compound for 6 days. Plating ability of the cells was assessed by plating
5 dilutions of each culture on plastic. Colony forming units were determined after growth for two to four days by counting colonies after staining the plates. Each sample was plated in duplicate and the colony number averaged. The results are set forth in **Figure 1**.

10 The third test for cell toxicity assessed growth rate in the presence of the test compound versus a control culture. Magic cells were grown over a period of 15 days with or without various dilutions of test compound. Aliquots were removed every other day
15 and the cells were counted in a hemocytometer. The results are set forth in **Figure 2** herein (see also Table 2 below).

EXAMPLE 5

Integrase Inhibition Assays

20 *Testing as Integrase Inhibitors.* Integrase overproduced in *E. coli* was purified as described above in accordance with Sherman et al, *supra*, and used in drug inhibition assays. The integrase preparation used for these studies was extremely pure and did not
25 contain any contaminating nuclease activity. Dilutions of the inhibitor compounds described in Example 1 above were mixed with substrate before the addition of enzyme. The Assay was the same as described in Example 2 and used the substrate described in Example 2. Each
30 assay was performed in duplicate. Radioactivity in bands on dried gels were quantitated with a Phosphor imager to assess drug effects on both the cleavage, nuclease, and ligation products. The IC_{50} values for integrase inhibition were calculated for both the
35 cleaving and joining parts of the integrase reaction after determining the % inhibition of the control reaction for a series of drug concentrations. These

-20-

results are shown in Table 2 below. Compounds A, B, C, and D had similar effects on integrase activity. Compound A, however, caused aggregation of the substrate starting at relatively low concentrations (10 μ M), complicating its assessment.

TABLE 2

Summary of IC₅₀s (μ M)

	Drug	Magic Cells		Integrase ¹	
		HIV-1	Toxicity	Cleavage	Strand Transfer
	A ²	48	>500	11	11.5
10	B	10-30	473.2	61	19
	C	16	237.9	64	22
	D	7	175.3	-	-
	E	16	65	>100	122

¹Activity was measured at the % cleavage of the substrate compared to control.

15 ²This drug caused aggregation of the substrate and also precipitated out of the culture medium at higher drug concentrations.

Although not wishing to be bound by any theory or explanation of the invention, these compounds are currently believed to bind in the double stranded DNA minor groove with an AT bias (see T. Fairley et al, *J. Med. Chem.* 36, 1746-1753 (1993)), and most likely inhibit integrase by preventing the binding of integrase to its recognition sequences at the long terminal repeat ("LTR") of the virus. This proposed mechanism is supported by the observation that both cleavage and integration are equally effected by the compounds. Either DNA sequence specificity and/or direct interactions with the integrase protein are also currently believed to be possibly involved in the compound mechanism. Since integrase functions as a multimer, K.S. Jones et al, *J. Biol. Chem.* 267, 16037-40 (1992), it is also possible that the DNA binding of the compounds somehow effects the multimer equilibrium.

-21-

The results indicate that DNA binding strength alone, however, is not the determining factor. Either DNA sequence specificity and/or direct interactions with the integrase protein are currently
5 also believed to be possibly involved. Since nucleosomes have been shown to be precisely positioned in the 5' LTR of HIV-1, A. Fesen et al., *Proc. Natl. Acad. Sci. USA* **90**, 2399-2403 (1993), such phasing could be one other way that the dicationic, groove binding
10 drugs interfere with integrase action.

EXAMPLE 6

HIV-1 Inhibition Assays

The magic cell assay described by Kimpton et al., *supra*, was used as described. This assay
15 identifies individual cells infected with HIV-1 by the expression of tat, which transactivates an endogenous copy of the HIV-1 LTR linked to the lac2 reporter gene after integration, inducing β -galactosidase expression when x-gal is added to the cell medium. Any cell with
20 integrated HIV-1 will turn blue. Thus, this assay provides a convenient way to determine the effect of HIV inhibitors at any early step up through the expression of tat, including the inhibition of integration.

25 The magic cells are plated in twelve-well plates one day prior to infection. The standard assay involves infection with approximately 200 infectious units of HIV-1. This gives an approximately 20 to 1 ratio of signal to background and sufficient numbers of
30 infectious events to quantitate dim effects. The cells are pretreated with drug for 4 hours prior to virus infection, and virus adsorption takes place for 1 hour. The cells are washed with plain medium, and then medium with inhibitor is placed back on the cells. Two days
35 later the cells are fixed after integration with x-gal, the indicator reagent for β -galactosidase production.

-22-

The number of β -galactosidase expressing cells are quantitated by light microscopy.

The results of comparisons of infectious units with or without various concentrations of the bis-benzimididazoles drugs in the magic cell assay are expressed as IC_{50} values and are compiled in Table 2 above. Note that the best anti-HIV compounds (B-D) were also the best integrase inhibitors, except for compound A, which could not be accurately measured due to its substrate aggregating properties.

In addition, the assessment of antiviral activity as measured by the standard reinfection assay was performed for compound B. The IC_{50} for compound B in this assay was 10-30 μ M. This confirms that the Magic cell assay is suitable for screening potential integrase inhibitors with good antiviral activity.

Another test of cellular effects by the compounds was a measure of plating efficiency of the cells after incubation with various concentrations of test agents. This will assess the actual killing of the cells by the drugs. The IC_{50} values are expected to be much higher in this test. The results indicate that the best anti-HIV compound, B, had no effect on the Magic cell plating efficiency at the concentrations at which anti-viral effects were observed.

The most sensitive test employed measurements of growth rates for the cells in the presence of the test compound. Although the anti-viral assay used was one that assessed early times of the viral life cycle, this test was carried out for 15 days, with counting of cell numbers in cultures performed every other day during this time. It is important to note that for up to two weeks, the concentration of Compound B that inhibited 50% of the viral integration events had no effect on the growth rate of the cell line. Higher concentrations, however, did effect the growth rate of the cells.

-23-


The foregoing examples are illustrative of the present invention, and are not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be

5 included therein.

-24-

THAT WHICH IS CLAIMED IS:

1. A method of inhibiting retroviral integrase in a subject in need of such treatment, comprising administering to said subject an effective retroviral inhibiting amount of a compound according to
- 5 Formula (I):



wherein:

- R_1 and R_2 are each independently selected from the group consisting of H or lower alkyl, or R_1 and R_2 together represent $-(CH_2)_m-$ wherein m is from two to
- 10 four;

R_3 is H or lower alkyl; and

X is C1 to C2 saturated or unsaturated alkyl containing up to one double bond;

- or a pharmaceutically acceptable salt
- 15 thereof.

2. A method according to claim 1, wherein R_1 , R_2 , and R_3 are H.

3. A method according to claim 1, wherein R_1 and R_2 together represent $-CH_2-CH_2-$ and R_3 is H.

- 20 4. A method according to claim 1, wherein X is $-(CH_2)_n-$ wherein n is from 1 to 2, which is unsubstituted or substituted from 1 to 2 times with lower alkyl, and which is saturated or unsaturated and contains up to one double bond.

-25-

5. A method according to claim 1, wherein X is selected from the group consisting of: $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, and any of the foregoing substituted from 1 to 2 times with lower alkyl.

5 6. A method according to claim 1, wherein said compound is selected from the group consisting of:
 (A) 1,2-bis[5-amidino-2-benzimidazolyl]ethene;
 (B) 1,2-bis[5-isopropylamidino-2-
10 benzimidazolyl]ethene;
 (C) bis[5-(2-imidazolyl)-2-benzimidazolyl] methane;
 (D) bis[5-amidino-2-benzimidazolyl] methane;
and the pharmaceutically acceptable salts thereof.

15 7. A method according to claim 1, wherein said compound of Formula I is bis[5-amidino-2-benzimidazolyl]methane or a pharmaceutically acceptable salt thereof.

 8. A method according to claim 1, wherein
20 said compound of Formula I is 1,2-bis[5-amidino-2-benzimidazolyl]ethane or a pharmaceutically acceptable salt thereof.


 9. A method according to claim 1, wherein said compound of Formula I is 1,2-bis[5-amidino-2-
25 benzimidazolyl]ethene or a pharmaceutically acceptable salt thereof.

 10. A method according to claim 1, wherein the subject is mammalian.

 11. A method according to claim 1, wherein
30 the subject is avian.

-26-

12. A method of combatting a retroviral infection in a subject in need of such treatment, comprising administering to said subject an effective retroviral infection combatting amount of a compound according to Formula (I):



wherein:

- R_1 and R_2 are each independently selected from the group consisting of H or lower alkyl, or R_1 and R_2 together represent $-(CH_2)_m-$ wherein m is from two to four;

R_3 is H or lower alkyl; and

X is C1 to C2 saturated or unsaturated alkyl containing up to one double bond;

- or a pharmaceutically acceptable salt thereof.

13. A method according to claim 12, wherein R_1 , R_2 , and R_3 are H.

14. A method according to claim 12, wherein R_1 and R_2 together represent $-CH_2-CH_2-$ and R_3 is H.

15. A method according to claim 12, wherein X is $-(CH_2)_n-$ wherein n is from 1 to 2, which is unsubstituted or substituted from 1 to 2 times with lower alkyl, and which is saturated or unsaturated and contains up to one double bond.

-27-

16. A method according to claim 12, wherein X is selected from the group consisting of $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, and any of the foregoing substituted from 1 to 2 times with lower alkyl.

5 17. A method according to claim 12, wherein said compound is selected from the group consisting of:
(A) 1,2-bis[5-amidino-2-benzimidazolyl]ethene;
(B) 1,2-bis[5-isopropylamidino-2-
10 benzimidazolyl]ethene;
(C) bis[5-(2-imidazolyl)-2-benzimidazolyl]methane;
(D) bis[5-amidino-2-benzimidazolyl] methane;
and the pharmaceutically acceptable salts thereof.

15 18. A method according to claim 12, wherein said compound of Formula I is bis[5-amidino-2-benzimidazolyl]methane or a pharmaceutically acceptable salt thereof.

19. A method according to claim 12, wherein
20 said compound of Formula I is 1,2-bis[5-amidino-2-benzimidazolyl]ethane or a pharmaceutically acceptable salt thereof.

20. A method according to claim 12, wherein said compound of Formula I is 1,2-bis[5-amidino-2-
25 benzimidazolyl]ethene or a pharmaceutically acceptable salt thereof.

21. A method according to claim 12, wherein the subject is mammalian.

22. A method according to claim 12, wherein
30 the subject is avian.




FIG. 1.




FIG. 2.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/00504

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/415 A61K31/155

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A,4 619 942 (TIDWELL ET AL) 28 October 1986 see the whole document ---	1-22
Y	J. CLIN. INVEST., vol. 82, no. 19, 1988 pages 2011-6, VONDERFECHT ET AL 'Protease Inhibitors Suppress the In Vitro and In Vivo Replication of Rotavirus' * see p.2015, Discussion * --- -/--	1-22

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 May 1995

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Uiber, P

INTERNATIONAL SEARCH REPORT

Inter- national Application No

PCT/US 95/00504

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. MED. CHEM., vol. 36, 1993 pages 1746-53, FAIRLEY ET AL 'Structure, DNA Minor Groove Binding, and Base Pair Specificity of Alkyl- and Aryl-Linked Bis(Amidinobenzimidazoles) and Bis(Amindinoindoles)' cited in the application see the whole document ---	1-22
A	AM. J. PATHOL., vol. 123, no. 1, 1986 pages 46-56, CRESSON ET AL 'In Vitro Inhibition of Human Sarcoma Cells' Invasive Ability by Bis(5-Amidino-2-Benzimidazolyl)methane - a Novel Esteroprotease Inhibitor' see the whole document ---	1-22
A	FR,A,2 081 401 (ABBOTT LAB.) 3 December 1971 see the whole document ---	1-22
A	J. OF VIROL., vol. 66, no. 12, 1992 pages 7414-9, LAFEMINA ET AL 'Requirement of Active Human Immunodeficiency Virus Type 1 Integrase Enzyme for Productive Infection of Human T-Lymphoid Cells ' cited in the application see the whole document ---	1-22
Y	PROC. NATL. ACAD. SCI. USA, vol. 90, 1993 pages 2399-403, FERSEN ET AL 'Inhibitors of Human Immunodeficiency Virus Integrase' cited in the application * p.2401, left hand col.; Table 1; Fig. 1 *	1-22
E	WO,A,95 08540 (THE WELLCOME FOUNDATION LTD.) 30 March 1995 * see p.11, 1.11-32; claims 2,3; p.9-10 * -----	1-22

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 95/00504

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4619942	28-10-86	US-A- 4397863	09-08-83
FR-A-2081401	03-12-71	CH-A- 535771	15-04-73
		DE-A- 2063856	01-07-71
WO-A-9508540	30-03-95	NONE	